

**This Page Is Inserted by IFW Operations
and is not a part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- **BLACK BORDERS**
- **TEXT CUT OFF AT TOP, BOTTOM OR SIDES**
- **FADED TEXT**
- **ILLEGIBLE TEXT**
- **SKEWED/SLANTED IMAGES**
- **COLORED PHOTOS**
- **BLACK OR VERY BLACK AND WHITE DARK PHOTOS**
- **GRAY SCALE DOCUMENTS**

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 83108583.2

(22) Date of filing: 31.08.83

(51) Int. Cl.³: C 07 D 487/04

C 07 D 231/04, C 07 D 237/04
 C 07 D 243/02, C 07 D 245/02
 A 01 N 43/90, A 01 N 47/38
 //(C07D487/04, 249/00, 237/00),
 (C07D487/04, 249/00, 231/00),
 (C07D487/04, 249/00, 243/00),
 (C07D487/04, 249/00, 245/00)

(30) Priority: 02.09.82 JP 151696/82
13.09.82 JP 158227/82

(43) Date of publication of application:
04.04.84 Bulletin 84/14

(84) Designated Contracting States:
AT BE CH DE FR GB IT LI NL SE

(71) Applicant: Nippon Kayaku Kabushiki Kaisha
New Kaijo Bldg. 2-1, Marunouchi 1-chome Chiyoda-ku
Tokyo 100(JP)

(72) Inventor: Kobayashi, Shinichi
4-8-24, Sengencho
Fuchu-shi Tokyo(JP)

(72) Inventor: Yanagi, Mikio
178-28, Oaza-Idogi Ageo-shi
Saitama-ken(JP)

(72) Inventor: Yamada, Osamu
473-5, Mukoyama
Ageo-shi Saitama-ken(JP)

(72) Inventor: Shida, Atsuhiko
11-6, 3-chome, Higashihonmachi
Koga-shi Ibaraki-ken(JP)


(72) Inventor: Futatsuya, Fumio
9-306, 710-50, Ohaza-higashierai
Ohmiya-shi Saitama-ken(JP)

(72) Inventor: Shimano, Shizuo
2-11-23, Kamicho
Ageo-shi Saitama-ken(JP)


(74) Representative: Türk, Dietmar, Dr. rer. nat. et al,
Redies, Redies, Türk & Gille Patentanwälte
Brucknerstrasse 20
D-4000 Düsseldorf 13(DE)

(54) A heterocyclic compound and its use as herbicide.

(57) A compound of the formula





wherein Z is



wherein R₄ is lower alkyl, X and Y are oxygen or sulfur
 and n is an integer of 3 to 6,
 R₁ is hydrogen or halogen, R₂ is halogen and R₃ is hydrogen
 or C₁ - C₆-alkyl which may have lower alkoxy; a herbicidal
 composition containing said compound as an effective
 component; a method for killing weeds using said com-
 pound; and processes for the production of said compound,
 are disclosed hereinafter.

This invention relates to a compound of the formula:



wherein R_4 is lower alkyl, X and Y are oxygen or sulfur and n is an integer of 3 to 6;

R_1 is hydrogen or halogen, R_2 is halogen and R_3 is hydrogen or $C_1 - C_8$ -alkyl which may have lower alkoxy; a herbicidal composition containing said compound as an effective component; a method for killing weeds using said compound; and processes for the production of said compound.

It is known that, among the compounds similar to the compound of the present invention, there are compounds having herbicidal activity (Japanese Patent Kokai No. 50-160429, 51-35435, 52-83687, 53-44587, 51-38425, 52-83552 and 53-40785).

The present inventors have found that the compound of the formula (1) has very high herbicidal activity. Furthermore, the present inventors have found that the compounds of the formula (1) have very low phytotoxicity against crops and, therefore, they become practical herbicide.


The compound of the formula (1) exhibits excellent

herbicidal effect in a paddy field at a low dosage not only against annual weeds such as barnyard grasses and broadleaf weeds, but also against perennial weeds such as mizugayatsuri, bulrush, water chestnut, needle spikerush and arrowhead. The compounds of the formula (1) also show a good herbicidal effect by both pre- and post-emergence treatments in an up-land, especially against broadleaf weeds as those of amaranth, goosefoot and buckwheat families at a low dosage.


Furthermore, the compound of the formula (1) is hardly phytotoxic to crops such as rice, wheat, oat, corn, soybean, cotton and sunflower.

The compound of the formula (1) of the present invention can be produced by the processes described in (A), (B), (C), (D) and (E) below.

(A):



The compound of the formula (2) wherein R_1 , R_2 , R_3 , R_4 , X , Y and n are as defined in formulae (3) and (4), which corresponds to a compound of the formula (1) wherein Z is




can be obtained as follows:

A compound of the formula (4) (for example, N^1, N^2 -trimethylene hydrazine, N^1, N^2 -hexamethylene hydrazine) can be reacted with a chlorocarbonic ester derivative of the formula (5) wherein R_4 is lower alkyl, Y is oxygen or sulfur in an organic solvent such as alcohols (methanol, ethanol) and the inert solvent defined after, at $0 - 50^\circ C$, preferably at $10 - 20^\circ C$ to produce an N^1, N^2 -alkylene- N^1 -alkoxycarbonylhydrazine derivative of the formula (6) wherein R_4 , Y are as defined in the formula (5), n is as defined in the formula (3), and the compound of the formula (6) can be reacted with a substituted phenylisocyanate or a substituted phenylthioisocyanate derivative of the formula (7) wherein R_1 , R_2 , R_3 and X are as defined in the formula (3) preferably in an inert solvent, at $0 - 100^\circ C$ more preferably at $5 - 50^\circ C$ for 20 min. to 3 hrs. to produce a compound of the formula (2).

(B):


The compound of the formula (3) wherein R_1 , R_2 , R_3 , X , Y and n are as defined above, which corresponds to the compound of the formula (1) wherein Z is




can be

obtained by cyclizing the compound of the formula (2) by heating preferably under reflux in the presence of a base at 60 - 150°C preferably at 80 - 120°C for 30 min. to 8 hrs.

(C):




The compound of the formula (11) wherein R_1 , R_2 , R_3 , X and n are as defined in the formula (1), which corresponds to the compound of the formula (1) wherein Z is $(\text{CH}_2)_n$ can be produced preferably by reacting a sodium salt



The compound of the formula (9) can be produced by cyclizing a compound of the formula (8) by heating, preferably under reflux, at 60 - 150°C, preferably at 80 - 120°C in the presence of a base in an inert solvent.


The compound of the formula (8) is obtained by reacting an N-alkyl carbonylhydrazine with the compound of the formula (7), preferably in an inert solvent, at $-10 - 100^{\circ}\text{C}$, more preferably at $-5 - 50^{\circ}\text{C}$.

(D):



Furthermore, the compounds of the formulae (13) and (14) wherein R_1 , R_2 , R_3 and n are as defined in the formula (3) can be obtained by reacting a urazol derivative of the formula (12) with a phosphorus polysulfide such as P_2S_5 under heating in an organic solvent such as benzene, toluene, xylene, preferably at $80 - 140^{\circ}\text{C}$, or reacting said urazol derivative with preferably B_2S_3 or SiS_2 in chloroform under reflux.

(E):




The former compound of the formula (2) also can be obtained by reacting the compound of the formula (4) with the compound of the formula (7) in an inert solvent preferably at $-10 - 30^\circ\text{C}$ to produce a compound of the formula (15) wherein R_1 , R_2 , R_3 , X and n are as defined in the formula (3), and then reacting thus obtained compound with a compound of the formula (5) in an inert solvent at $0 - 50^\circ\text{C}$.

The compounds of the formulae (4) and (6) which are starting materials of the compound of the formula (1) can be obtained by the known method (Japanese Patent Kokai No. 51-65757).

The compound of the formula (7) can be obtained by reacting a substituted aniline derivative of the formula (16) wherein R_1 , R_2 , R_3 are as defined in the formula (3) with

phosgene or thiophosgene of the formula (17) or ClCOOCCl_3 .



The above reaction can be conducted in the absence of a solvent, but preferably it is conducted in an inert solvent, generally at -20°C - boiling point of the solvent, preferably at 0°C - 120°C .

The reaction is conducted preferably under atmospheric pressure, but it can be conducted under pressure or reduced pressure.

As examples of the bases used in the present invention, there can be mentioned alkali metal salts such as sodium hydroxide or potassium hydroxide; alkaline earth metal salts such as calcium hydroxide or magnesium hydroxide and barium hydroxide; alcoholates of alkali metal such as sodium alcoholates or potassium alcoholates; trialkylamines; pyridine; picoline, and usually the bases can be used in catalytic amount.

As examples of the inert solvents, there can be mentioned aromatic hydrocarbons such as benzene, toluene, xylenes, chlorobenzenes; aliphatic hydrocarbons such as n-hexane, n-heptane or


petroleum ether; alicyclic hydrocarbon such as cyclohexane; halogenated hydrocarbons such as chloroform, carbontetrachloride or perchloroethylene; ketones such as acetone or methylethylketone; ethers such as ethylether, tetrahydrofuran or dioxane; alcohols such as methanol or ethanol; esters such as ethylacetate or butylacetate; amides such as N,N-dimethylformamide or N,N-diethylformamide; or water.

In the formula (1), as examples of halogens there can be mentioned chloro, bromo or fluoro. As examples of lower alkoxys there can be mentioned methoxy or ethoxy.


As examples of $C_1 - C_8$ -alkyls, there can be mentioned methyl, ethyl, n-propyl, iso-propyl, n-butyl, s-butyl, iso-butyl, tert-butyl, n-pentyl, n-hexyl, n-heptyl, n-octyl, 2-ethylhexyl or 1,3-dimethylbutyl.

In the formula (4), as examples of lower alkyls of R_4 , there can be mentioned $C_1 - C_4$ alkyls such as methyl, ethyl, n-propyl, iso-propyl, n-butyl or sec-butyl.

As preferable compounds in the present invention, there can be mentioned those of the formula (1) wherein (a) Z is




R_1 is chloro or fluoro, R_2 is chloro or bromo, R_3 is $C_2 - C_4$ -alkyl, (b) Z is



are oxygen or sulfur), R_1 is chlor or fluoro, R_2 is chloro or bromo and R_3 is $C_2 - C_4$ -alkyl.

As most preferable compounds in the present invention, there can be mentioned



The present invention will be illustrated in the following examples.

Synthesis example 1:

1,2-trimethylene-4-(2-fluoro-4-chloro-5-ethoxycarbonylphenyl)urazol (Compound No.1).

1.44 g (0.01 mol) of 1,2-trimethylenehydrazine-1-carboxylic acid ethylester were dissolved in 30 ml of toluene and 2.3 g (0.01 mol) of 2-fluoro-4-chloro-5-ethoxycarbonylphenylisocyanate were added to the solution, and after the mixture were stirred for 1 hr. at room temperature, 2 ml of N,N-dimethylformamide and 0.3 g of sodium hydroxide were added thereto and the whole was stirred for 15 min. at 100 - 110°C. After cooling, the reaction mixture was washed with water, dried over anhydrous Na_2SO_4 and concentrated to obtain crystals. Upon recrystallization from ethanol, 1.21 g of pale brown crystals were obtained.

Yield: 37.2%, M.p.: 156 - 158°C.

Elementary analysis: $C_{14}H_{18}FClN_3O_4$

calculated: C: 49.20, H: 3.83, N: 12.29

found: C: 49.50, H: 3.88, N: 12.32

Synthesis example 2:

1,2-tetramethylene-4-(2-fluoro-4-bromo-5-isopropoxy-carbonylphenyl)urazol (Compound No.13).

1.6 g (0.01 mol) of 1,2-tetramethylenedihydrazine-1-carboxylic acid ethylester were dissolved in 30 ml of toluene and 3 g (0.01 mol) of 2-fluoro-4-bromo-5-isopropoxycarbonylphenylisocyanate were added to the solution, and after the mixture was stirred for 1 hr. at room temperature, 2 ml of N,N-dimethylformamide and 0.3 g of sodium hydroxide were added thereto and the whole was heated at 80°C for 30 min. After cooling, the reaction mixture was washed with water, dried over anhydrous Na_2SO_4 and concentrated to obtain crystals. Upon recrystallization from toluene, 27.1 g of white crystals were obtained. Yield: 65%, M.P.: 161 - 162°C.

Elementary analysis: $C_{16}H_{17}FBrN_3O_4$

Calculated: C: 46.39, H: 4.13, N: 10.14

Found: C: 46.68, H: 4.14, N: 10.30

Synthesis example 3:

1,2-tetramethylene-4-(4-chloro-3-methoxycarbonylphenyl)-thiourazol (Compound No. 14).

1.6 g (0.1 mol) of 1,2-tetramethylenedihydrazine-1-carboxylic acid ethylester were dissolved in 40 ml of xylene and

when 2.2 g (0.01 mol) of 4-chloro-3-methoxycarbonylphenylisothiocyanate were added to the solution, white crystals were precipitated. After precipitating, the mixture was stirred at room temperature for 1 hr., 200 mg of hydrous Na_2SO_4 were added thereto and the whole was heated under reflux for 3 hrs. After cooling, the reaction mixture was washed with water, dried over anhydrous Na_2SO_4 and concentrated to obtain crystals. Upon recrystallization from the mixture of n-hexane and toluene, 2.2 g of pale yellow crystals were obtained.

Yield: 64%, M.P.: 126 - 128°C

Elementary analysis: $\text{C}_{14}\text{H}_{14}\text{ClN}_3\text{O}_3\text{S}$

Calculated: C: 49.48, H: 4.15, N: 12.36

Found: C: 49.41, H: 4.19, N: 12.08

Synthesis example 4:

1,2-tetramethylene-4-(4-chloro-3-isopropoxycarbonylphenyl)dithiourazol (Compound No.19).

3.6 g of (0.01 mol) 1,2-tetramethylene-4-(4-chloro-3-isopropoxycarbonylphenyl)thiourazol and 2.2 g (0.01 mol) of diphosphorus pentasulfide were added to 60 ml of xylene and heated under stirring at 100 - 110°C for 9 hrs.

After cooling, the produced precipitates were removed by filtration and the filtrate was washed with water, dried over anhydrous Na_2SO_4 and then the oil was obtained by concentration. Thus obtained oil was purified by the column chromatography (silica gel) to obtain 1.75 g of yellow crystals. Yield: 40%, M.p.: 161 - 163°C.

Elementary analysis: $C_{16}H_{18}ClN_3O_2S_2$

Calculated: C: 50.05, H: 4.72, N: 10.94

Found: C: 50.21, H: 4.77, N: 10.69

Synthesis example 5:

1,2-tetramethylene-4-(2-fluoro-4-chloro-5-sec-butoxycarbonylphenyl)thiourazol (Compound No. 26).

8.3 g (0.0175 mol) of 1,2-tetramethylene-1-carbo-iso-butoxy-2-(fluoro-4-chloro-5-sec-butoxycarbonyl phenylthiocarbamoyl)-hydrazine were dissolved in 100 ml of toluene, 3 ml of N,N-dimethylformamide and 0.5 g of sodium hydroxide were added to the solution and the mixture was heated at 100 - 110°C for 30 min. After cooling, the reaction mixture was washed with water, dried over anhydrous Na_2SO_4 , and concentrated to obtain crystals. Upon recrystallization from the mixture of n-hexane and toluene, 5 g of white crystals were obtained. Yield: 71.4%, M.p.: 131.5 - 132.5°C.

Elementary analysis: $C_{17}H_{19}FClN_3O_3S$

Calculated: C: 51.06, H: 4.78, N: 10.50

Found: C: 51.29, H: 4.82, N: 10.45

Synthesis example 6:

1,2-pentamethylene-4-(2-fluoro-4-chloro-5-isopropoxy-carbonylphenyl)urazol (Compound No. 30).

2.9 g (0.0067 mol) of 1,2-pentamethylene-1-carboethoxy-2-(2-fluoro-4-chloro-5-isopropoxycarbonylphenylcarbamoyl)hydrazine were dissolved in 70 ml of toluene, 2 ml of N,N-dimethylformamide and 0.3 g of sodium hydroxide were added to the solution

0104484

and the mixture was heated under reflux for 30 min. After cooling, the reaction mixture was washed with water, dried over anhydrous Na_2SO_4 and concentrated to obtain crystals. Upon recrystallization from the mixture of n-hexane and toluene, 2.5 g of white crystals were obtained. Yield: 97.2%, M.p.: 105.5 - 106°C.

Elementary analysis: $\text{C}_{17}\text{H}_{19}\text{FClN}_3\text{O}_4$

Calculated: C: 53.20, H: 4.98, N: 10.94

Found: C: 53.43, H: 4.95, N: 11.18

Synthesis example 7:

1,2-pentamethylene-4-(4-bromo-3-isopropoxycarbonylphenyl)thiourazol (Compound No. 33).

1.5 g (0.0087 mol) of 1,2-pentamethylenedihydrazine-1-carboxylic acid ethylester were dissolved into 40 ml of xylene, 2.6 g (0.0087 mol) of 4-bromo-3-isopropoxycarbonylphenylisothiocyanate were added to the solution and the mixture was stirred for 1 hr., 300 mg of anhydrous Na_2SO_4 were added to the mixture and then the whole was heated under reflux for 12 hrs. After cooling, the reaction mixture was washed with water, dried over anhydrous Na_2SO_4 and concentrated to obtain the oil. Thus obtained oil was purified by the column chromatography (silica gel) to obtain 1.5 g of yellow oil. Yield: 40.2%, reflectance: n_D^{25} 1.5959


Elementary analysis: $\text{C}_{17}\text{H}_{20}\text{BrN}_3\text{O}_3\text{S}$

Calculated: C: 47.89, H: 4.72, N: 9.85

Found: C: 48.15, H: 4.69, N: 9.92

The following compounds were obtained by the above processes.

Table 1



Compound No.	n	X	Y	R_1	R_2	R_3	M.p.(°C) or refractive index	Appearance
1	3	O	O	F	Cl	C_2H_5	156-158	pale brown crystal
2	3	O	S	F	Cl	C_2H_5	amorphous	"
3	3	O	S	Cl	Cl	C_2H_5	132-134	pale yellow crystal
4	3	O	S	H	Cl	C_3H_7 (i)	169-171	"
5	3	O	S	F	Cl	C_3H_7 (i)	178-180	pale brown crystal
6	3	O	S	H	Br	C_4H_9 (sec)	111-113	pale yellow crystal
7	4	O	O	F	Cl	CH_3	131-133	white crystal
8	4	O	O	F	Cl	C_2H_5	121-123	"
9	4	O	O	H	Cl	C_3H_7 (i)	100-100.5	"
10	4	O	O	H	Br	C_3H_7 (i)	128.5-130	"
11	4	O	O	Cl	Cl	C_3H_7 (i)	140-142	"
12	4	O	O	F	Cl	C_3H_7 (i)	147-148	"
13	4	O	O	F	Br	C_3H_7 (i)	161-162	"
14	4	O	S	H	Cl	CH_3	126-128	pale yellow crystal
15	4	O	S	H	Br	C_2H_5	105-107	yellow crystal

0104484

Compound No.	n	X	Y	R ₁	R ₂	R ₃	M.p.(°C) or refractive index	Appearance
16	4	O	S	H	Cl	C ₂ H ₅	104-106	pale brown crystal
17	4	O	S	H	Cl	C ₃ H ₇ (n)	72-74	yellow crystal
18	4	O	S	H	Cl	C ₃ H ₇ (i)	120-127	pale yellow crystal
19	4	S	S	H	Cl	C ₃ H ₇ (i)	161-163	yellow crystal
20	4	O	S	H	Br	C ₃ H ₇ (i)	133.5-135.5	white crystal
21	4	O	S	Cl	Cl	C ₂ H ₅	134-135.5	"
22	4	O	S	F	Cl	CH ₃	188-190	"
23	4	O	S	F	Cl	C ₂ H ₅	130-132	pale yellow crystal
24	4	O	S	F	Cl	C ₃ H ₇ (i)	132-133.5	white crystal
25	4	O	S	F	Br	C ₃ H ₇ (i)	147-148.5	pale brown crystal
26	4	O	S	F	Cl	C ₄ H ₉ (sec)	131.5-132.5	white crystal
27	4	O	S	H	Cl	C ₄ H ₉ (sec)	88-90	pale brown crystal
28	4	O	S	H	Br	C ₄ H ₉ (sec)	88-90	pale yellow crystal
29	5	O	O	H	Br	C ₃ H ₇ (i)	111-112	white crystal
30	5	O	O	F	Cl	C ₃ H ₇ (i)	105.5-106	"
31	5	O	S	H	Cl	CH ₃	155-158	pale yellow crystal
32	5	O	S	H	Cl	C ₃ H ₇ (i)	n _D ²⁵ 1.5762	yellow oil
33	5	O	S	H	Br	C ₃ H ₇ (i)	n _D ²⁵ 1.5959	"
34	5	O	S	F	Cl	C ₃ H ₇ (i)	142-143.5	white crystal
35	5	O	S	F	Br	C ₃ H ₇ (i)	171-173	brown crystal
36	4	O	S	F	Cl	$\begin{array}{c} \text{CH}-\text{CH}_2-\text{CH}-\text{CH}_3 \\ \quad \quad \\ \text{CH}_3 \quad \text{CH}_3 \end{array}$	n _D ²⁵ 1.5620	pale yellow oil
37	4	O	S	F	Cl	CH ₂ CH ₂ OCH ₃	83-85	white crystal
38	4	O	S	F	Cl	$\begin{array}{c} \text{CH}-\text{CH}(\text{CH}_2)_3-\text{CH}_3 \\ \quad \quad \\ \text{CH}_2 \quad \text{CH}_3 \end{array}$	n _D ²⁵ 1.5551	pale yellow oil

Synthesis example 8:

1,2-trimethylene-1-ethoxycarbonyl-2-(2-fluoro-4-chloro-5-ethoxycarbonylphenylcarbamoyl)hydrazine (Compound No. 40).

2.46 g (0.017 mol) of 1,2-trimethylenehydrazine were dissolved in 30 ml of toluene and 4.14 g (0.017 mol) of 2-fluoro-4-chloro-5-ethoxycarbonylphenylisocyanate were added to the solution and the whole was stirred at room temperature for 1 hr. The reaction mixture was washed with water, dried over anhydrous Na_2SO_4 and concentrated to obtain the crystal. The crystals were purified by column chromatography (silica gel, developing solvent: hexane : ethylacetate) to produce 4.29 g of white crystals, Yield: 65%, M.p.: 82-83°C.

Elementary analysis: $\text{C}_{16}\text{H}_{19}\text{FClN}_3\text{O}_5$

Calculated: C:49.55, H: 4.93, N: 10.83

Found: C:49.75, H: 4.94, N: 10.91

Even if 1,2-trimethylenehydrazine-1-carboxylic acid hydrochloride was used instead of 1,2-trimethylenehydrazine-1-carboxylic acid, the above compound (Compound No. 40) was obtained by using triethylamine.

Synthesis example 9:

1,2-trimethylene-1-carboethoxy-2-(4-bromo-3-sec-butoxycarbonylphenylthiocarbamoyl)hydrazine (Compound No. 45).

1.44 g (0.01 mol) of 1,2-trimethylenehydrazine-1-carboxylic acid ethylester were dissolved in 30 ml of benzene and 3.1 g (0.01 mol) of 4-bromo-3-sec-butoxycarbonylphenylisothiocyanate were added to the solution and the mixture was

0104484

stirred for 1 hr. at room temperature. The reaction mixture was concentrated to obtain the crystals. Upon r crystallization from the mixture of hexane and toluene, 3.9 g of white crystals were obtained. Yield: 85%, M.p.: 119 - 120.5°C.

Elementary analysis: $C_{18}H_{24}BrN_3O_4S$

Calculated: C: 47.16, H: 5.27, N: 9.16

Found: C: 47.08, H: 5.53, N: 9.26

The following compounds were obtained by the above processes.




Table 2

Compound No.	n	X	R ₁	R ₂	R ₃	R ₄	M.p.(°C) or refractive index	Appearance
39	3	O	F	Cl	CH ₃	C ₂ H ₅	103-105	white crystal
40	3	O	F	Cl	C ₂ H ₅	"	82-83	"
41	3	O	F	Cl	C ₃ H ₇ (i)	"	102-105	"
42	3	S	H	Cl	CH ₃	"	127-128	pale yellow crystal
43	3	S	H	Cl	C ₃ H ₇ (i)	"	108-109	"
44	3	S	F	Cl	C ₃ H ₇ (i)	"	87-88.5	pale brown crystal
45	3	S	H	Br	C ₄ H ₉ (sec)	"	119-120.5	white crystal
46	3	S	Cl	Cl	C ₂ H ₅	"	102-104	"
47	3	S	F	Cl	C ₂ H ₅	"	97-98.5	pale brown crystal
48	4	O	H	Cl	C ₃ H ₇ (i)	"	119-122	"
49	4	O	F	Cl	CH ₃	"	126.5-127	white crystal
50	4	O	F	Cl	C ₂ H ₅	"	87-88	"
51	4	O	F	Cl	C ₃ H ₇ (i)	"	n _D ²⁵ 1.5249	yellow oil
52	4	O	H	Br	C ₃ H ₇ (i)	"	125-126	white crystal
53	4	O	Cl	Cl	C ₃ H ₇ (i)	"	69-71	"
54	4	O	F	Br	C ₃ H ₇ (i)	"	74-75	"
55	4	O	F	Cl	C ₃ H ₇ (i)	C ₄ H ₉ (i)	n _D ²⁵ 1.5161	pale yellow oil
56	4	O	F	Br	C ₃ H ₇ (i)	"	n _D ²⁵ 1.5216	"
57	4	S	H	Cl	CH ₃	CH ₃	181.5-183.5	pale yellow crystal
58	4	S	H	Cl	CH ₃	C ₂ H ₅	144-146	white crystal
59	4	S	H	Cl	CH ₃	C ₄ H ₉ (i)	108-109.5	pale yellow crystal
60	4	S	H	Cl	C ₂ H ₅	C ₂ H ₅	141-143	pale brown crystal

Compound N .	n	X	R ₁	R ₂	R ₃	R ₄	M.p.(°C) refractive index	Appearance
61	4	S	Cl	Cl	C ₂ H ₅	C ₂ H ₅	88-90	white crystal
62	4	S	F	Cl	"	"	129-130.5	"
63	4	S	H	Cl	C ₃ H ₇ (i)	CH ₃	100-102	pale brown crystal
64	4	S	F	Br	C ₃ H ₇ (i)	CH ₃	amorphous	dark yellow
65	4	S	H	Cl	C ₃ H ₇ (n)	C ₂ H ₅	132-134	pale brown crystal
66	4	S	H	Cl	C ₃ H ₇ (i)	"	124-126	pale yellow crystal
67	4	S	H	Br	C ₃ H ₇ (i)	"	121-123	"
68	4	S	F	Br	C ₃ H ₇ (i)	"	n _D ²⁵ 1.5514	brown oil
69	4	S	H	Cl	C ₃ H ₇ (i)	C ₄ H ₉ (i)	123-124	white crystal
70	4	S	H	Br	C ₃ H ₇ (i)	"	124-126	pale yellow crystal
71	4	S	F	Br	C ₃ H ₇ (i)	"	n _D ²⁵ 1.5421	brown oil
72	4	S	F	Cl	"	"	n _D ²⁵ 1.5448	yellow oil
73	4	S	H	Br	C ₄ H ₉ (sec)	CH ₃	80-82	white crystal
74	4	S	H	Cl	"	C ₂ H ₅	101-103	"
75	4	S	H	Br	"	"	100-101.5	pale yellow crystal
76	4	S	F	Cl	"	C ₄ H ₉ (i)	n _D ²⁵ 1.5450	pale brown oil
77	5	O	F	Cl	C ₂ H ₅	C ₂ H ₅	81-83	pale brown crystal
78	5	O	H	Br	C ₃ H ₇ (i)	"	119-122	white crystal
79	5	O	Cl	Cl	"	"	68-71	"
80	5	O	F	Cl	"	"	80-81	"
81	5	S	F	Cl	C ₂ H ₅	"	amorphous	pale brown
82	5	S	H	Cl	C ₃ H ₇ (i)	"	121-122.5	white crystal
83	5	S	H	Br	"	"	114.5-116	"

Compound No.	n	X	R ₁	R ₂	R ₃	R ₄	M.p. (°C) refractive index	Appearance
84	5	S	Cl	Cl	C ₃ H ₇ (i)	C ₂ H ₅	108.5-110.5	white crystal
85	5	S	F	Cl	"	"	amorphous	pale yellow
86	5	S	F	Br	"	"	99-101	pale brown crystal
87	4	S	F	Cl	$\begin{array}{c} \text{CHCH}_2\text{CHCH}_3 \\ \quad \\ \text{CH}_3 \quad \text{CH}_3 \end{array}$	"	n _D ²⁵ 1.5480	pale yellow oil
88	4	S	F	Cl	CH ₂ CH ₂ OCH ₃	"	n _D ²⁵ 1.5547	"
89	4	S	F	Cl	$\begin{array}{c} \text{CH}_2\text{CH}(\text{CH}_2)_3\text{CH}_3 \\ \\ \text{C}_2\text{H}_5 \end{array}$	C ₂ H ₅	n _D ²⁵ 1.5370	"
90	4	S	F	Cl	C ₃ H ₇ (i)	"	n _D ²⁵ 1.5500	"

The compound of the formula (7) and the compound of the formula (2) were obtained as follows.

1. Synthesis of 4-chloro-2-fluoro-5-isopropoxycarbonylphenyl-isocyanate:

5.4 ml (0.045 mol) of trichloromethylchloroformate were added to 30 ml of ethylacetate and the solution was cooled to 0°C.

Then the mixture of 13.9 g (0.06 mol) of 4-chloro-2-fluoro-5-isopropoxycarbonylaniline and 25 ml of ethylacetate were added dropwise to the above solution over the period of 15 min. and the whole was stirred for 1 hr. at 0 - 10°C, thereafter heated under reflux for 2 hrs. After concentrating the reaction mixture, 15.45 g of white crystals were obtained. Yield: 99.9%, M.p.: 41 - 43°C.

2. Synthesis of 2,4-dichloro-5-ethoxycarbonylphenylisothiocyanate

2.34 g (0.1 mol) of 2,4-dichloro-5-ethoxycarbonylaniline were dissolved in 80 ml of chloroform and after the solution was cooled to below 10°C, 14.9 g (0.13 mol) of thiophosgene were added to the solution dropwise and after stirred for 2 hrs. at room temperature, the whole was heated under reflux for 3 hrs., followed by concentration to obtain 27 g of pale brown crystals. Yield: 97.7%, M.p.: 45 - 46°C.

The herbicidal composition of the present invention can be used either alone or in the form of a formulation according to the purpose of its use. To promote or secure the effect, it is mixed with adjuvants to make formulations such as dust, micro granule, granule, wettable powder, flowable suspension concentrates and emulsion by means of usual procedures. These formulations are used, at the time of practical application, in the form as they are or diluted with water to desired concentration.

Those adjuvants mentioned above include carriers (diluent), extending agents, emulsifiers, wetting agents, dispersing agents, fixing agents and disintegrators.

As liquid carriers there can be used water, aromatic hydrocarbons such as toluene and xylene, alcohols such as methanol, butanol and glycol, ketones such as acetone, amides such as dimethylformamide, sulfoxides such as dimethylsulfoxide, methylnaphthalene, cyclohexane, animal and vegetable oils, fatty acids and their esters, etc. As solid carriers are used

clay, kaolin, talc, diatomaceous earth, silica, calcium carbonate, montmorillonite, bentonite, feldspar, quartz, alumina, sawdust, etc.

As emulsifiers or dispersing agents surfactants are generally used. They include anionic, cationic, nonionic and amphoteric surfactants such as sodium salts of sulfated higher alcohol, stearyltrimethylammonium chloride, polyoxyethylenealkylphenylether and lauryl betaine. Wetting agents include sodium alkyl-naphthalene sulfonate and ammonium polyoxyethylenealkylphenylether sulfate, fixing agents include polyvinyl alcohol, polyvinyl acetate and CMC, and disintegrators include sodium lignin sulfonate.

Any type of said formulations can not only used alone, but also may be mixed with fungicides, insecticides, plant growth regulators, acaricides, soil modifying agents or nematocides and further can be used in combination with fertilizers or other herbicides.

The content of a compound (active ingredient) of the present invention in the formulations varies with types of formulation, methods of application and other conditions, but generally it is 0.5 to 95 weight %, preferably 2 to 50 weight %, while the content of adjuvants is 5 to 99.5 weight %, preferably 50 to 98 weight %, though sometimes the compound can be used alone.

To be more precise, a preferable range of the content is shown as under.

	Compound (weight %)	Adjuvant (weight %)
Dust	0.5 - 10	90 - 99.5
Emulsion	20 - 80	20 - 80
Wettable powder	20 - 80	20 - 80
Granule and micro granule	0.5 - 20	80 - 99.5
Flowable suspension concentrate	20 - 80	20 - 80

The formulations of the present invention can be directly applied to weeds or locus thereof.

A quantity to use of the formulations is different with kinds of the active ingredient and places of application, but generally it is within the range of 1 to 100 g, preferably 3 to 75 g, of the compound per are.

Detailed explanation will be made below on examples of formulations of the present invention and there the word "part" means part by weight.

Formulation Example 1 : Emulsion

35 parts of a mixture (1 : 1) of xylene and methyl-naphthalene are added to 50 parts of Compound No. 2 to dissolve and the solution is further mixed with 15 parts of a mixture (8 : 2) of polyoxyethylenealkylphenylether and calcium alkylbenzenesulfonate to obtain an emulsion. It is diluted with water to use in a concentration of 0.01 to 1 %.

Formulation Example 2 : Dust

5 parts of Compound No. 24 are mixed with 95 parts of clay and pulverized to obtain a dust. It is directly used for dusting.

Formulation Example 3 : Wettable powder

50 parts of Compound No. 25 are mixed with 10 parts of diatomaceous earth and 32 parts of kaolin and further uniformly blended with 8 parts of a mixture of sodium laurylsulfate and sodium 2,2'-dinaphthylmethanesulfonate, and finely pulverized to obtain a wettable powder. It is used in the form of a suspension by diluting to a concentration of 0.06 to 1 %.

Formulation Example 4 : Granule

5 parts of a fine dust of Compound No. 24 are extended for coating on 94.5 parts of grains (16 to 32 mesh) of silica to obtain a granule, by using a methanol solution of 0.5 parts of polyvinyl polyacetate as the binding agent in a proper mixer. The granule is scattered directly in up-land field and a paddy field.

Formulation Example 5 : Flowable suspension concentrates

40 parts of a fine powder of Compound 24, 10 parts of ethyleneglycolmonobutylether, 10 parts of a surfactant (mixture of trioxyalkylether, polyoxyethylenenonylphenylether and sodium alkylarylsulfonate), 3 parts of colloidal aluminium silicate hydrate and 22 parts of water are uniformly mixed and further blended under stirring in a homomixer for 20 minutes to obtain a flowable. It is diluted with water for use in a concentration of 0.02 to 1 %.

The excellent herbicidal activity of a compound of the present invention will be illustrated in the following test examples.

Each test was carried out on 2-replication system and the test results are given in the average value.

Test Example 1 : Pre-emergence treatment in flooded condition

A fixed amount of paddy field soil was filled in each Wagner pot sized 1/5,000 are to provide a condition similar to a paddy field and there was sown a fixed amount of seeds of barnyard grass, monochoria, toothcup, false pimpernel, water wort and umbrella plant.

In addition tubers of arrowhead were buried 1 cm under the surface of soil at the rate of 3 pieces per pot and the pot was flooded with water 3 cm deep. Then the pot was applied with a diluted solution of the compound of the present invention at a rate of 6.25 to 25 g of the compound of the present invention per are.

Thirty days after the treatment the herbicidal activity was observed. The test results were classified on the following basis as shown in Table 3.

Herbicidal activity index:

5	Complete weeding
4	up to 80% "
3	" 60% "
2	" 40% "
1	" 20% "
0	no effect

Table 3

Test Example 1 : Pre-emergence treatment under
flooded condition

Com- pound No.	Dosage g/a	Herbicidal activity			
		Barnyard grass	Broadleaf (1)	Umbrella- sedge (2)	Arrow-head
1	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
2	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
4	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
6	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
12	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
18	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
23	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
24	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
26	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
28	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5

0104484

Compound No.	Dosage .g/a	Herbicidal activity			
		Barnyard grass	Broadleaf (1)	Umbrella-sedge (2)	Arrow-head
30	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
35	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
40	25	5	5	5	4
	12.5	4	5	5	3
	6.25	3	5	5	2
41	25	5	5	5	3
	12.5	5	5	5	2.5
	6.25	3	5	5	2
42	25	5	5	5	5
	12.5	5	5	5	5
	6.25	4.5	5	5	5
43	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
44	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
45	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
47	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
48	25	5	5	5	4
	12.5	5	5	5	3
	6.25	2	5	5	2
50	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5


Compound No.	Dosage g/a	Herbicidal activity			
		Barnyard grass	Broadleaf (1)	Umbrella-sedge (2)	Arrow-head
51	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
52	25	5	5	5	5
	12.5	3	5	5	3
	6.25	2	5	5	2
54	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
56	25	5	5	5	5
	12.5	5	5	5	4
	6.25	3	5	5	3
57	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
59	25	5	5	5	5
	12.5	5	5	5	5
	6.25	3	5	5	5
60	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
61	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
62	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
63	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
64	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5

Compound No.	Dosage g/a	Herbicidal activity			
		Barnyard grass	Broadleaf (1)	Umbrella-sedge (2)	Arrow-head
66	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
68	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
71	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
72	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
74	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
75	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
76	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
81	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
82	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
83	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
85	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5

Compound No.	Dosage g/a	Herbicidal activity			
		Barnyard grass	Broadleaf (1)	Umbrella-sedge (2)	Arrow-head
86	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
88	25	5	5	5	5
	12.5	5	5	5	5
	6.25	5	5	5	5
Control A	25	3	4	3	0
	12.5	1	2	0	0

Remarks (1) Broadleaf: mixture of barnyard grass, toothcup, false pimpernel, water wort

(2) Umbrella-sedge: Umbrella plant



Test Example 2: Post-emergence treatment in flooded condition

A fixed amount of paddy field soil was filled in each Wagner pot sized 1/5,000 are to provide a condition similar to a paddy field and there was sown a fixed amount of seeds of barnyard grass, monochoria, toothcup, false pimpernel, water wort and umbrella plant.

In addition tubers of arrowhead were buried 1 cm under the surface of soil at the rate of 3 pieces per pot, three 2.5-leaf stage rice seedlings (variety : Nihonbare) were transplanted

from a nursery, the pot was flooded with water 3 cm deep and then placed in a greenhouse.

When the weeds grew to reach 2 to 3-leaf stage, a diluted solution of the compound of the present invention, was applied to the flood at a rate of 6.25, 12.5, 50 g of the compound of the present invention per are.

After 30 days from the treatment with the diluted solution, the phytotoxicity against paddy rice and the herbicidal activity were observed and obtained the results as shown in Table 4. The classification basis of the herbicidal activity is the same with Test Example 1, and that of the phytotoxicity is as follows:

- no damage
- + slight damage
- ++ some damage
- +++ moderate damage
- ++++ heavy damage
- x complete death

Table 4

Test Example 2: Post-emergence treatment under flooded condition

Compound No.	Dosage g/a	Herbicidal activity				Phytotoxicity against paddy rice
		Barnyard grass	Broad-leaf(1)	Umbrella sedge(2)	Arrow-head	
3	25	5	5	5	5	-
	12.5	4	5	5	5	-
10	25	5	5	5	5	-
	12.5	5	5	5	5	-
11	25	5	5	5	5	+
	12.5	5	5	5	5	-
14	25	5	5	5	5	+
	12.5	4.5	5	5	5	+
15	25	5	5	5	5	+
	12.5	5	5	5	5	+
19	25	5	5	5	5	-
	12.5	5	5	5	5	-
22	25	5	5	5	5	-
	12.5	4	5	5	5	-
27	25	5	5	5	5	+
	12.5	5	5	5	5	-
29	25	5	5	5	5	-
	12.5	5	5	5	5	-
31	50	5	5	5	5	+
	25	5	5	5	5	+
33	50	5	5	5	5	+
	25	5	5	5	5	+
34	25	5	5	5	5	++
	12.5	5	5	5	5	+
35	25	5	5	5	5	++
	12.5	5	5	5	5	+

Compound No.	Dosage g/a	Herbicidal activity				Phytotoxicity against paddy rice
		Barnyard grass	Broad-leaf(1)	Umbrella sedge(2)	Arrow-head	
39	25	3	5	5	3	-
	12.5	2.5	5	5	2	-
	6.25	2	5	5	2	-
45	25	5	5	5	4.5	-
	12.5	5	5	5	4	-
	6.25	2	5	5	3	-
46	25	5	5	5	5	-
	12.5	5	5	5	5	-
	6.25	5	5	5	4.5	-
47	25	5	5	5	5	++
	12.5	5	5	5	5	+
	6.25	5	5	5	5	-
49	25	3	5	5	3.5	-
	12.5	2	5	5	3	-
	6.25	2	5	5	2	-
51	25	5	5	5	5	-
	12.5	4.5	5	5	4	-
	6.25	4	5	5	3	-
52	25	5	5	5	4	-
	12.5	4	5	5	3	-
	6.25	3	5	5	2	-
54	25	5	5	5	5	-
	12.5	5	5	5	5	-
	6.25	3	5	5	4	-
57	25	5	5	5	5	+
	12.5	5	5	5	5	+
	6.25	4.5	5	5	5	-
59	25	5	5	5	5	++
	12.5	5	5	5	5	+
	6.25	5	5	5	5	+
60	25	5	5	5	5	+
	12.5	5	5	5	5	+
	6.25	5	5	5	5	+

Compound No.	Dosage g/a	Herbicidal activity				Phytotoxicity against paddy rice
		Barnyard grass	Broad-leaf(1)	Umbrella sedge(2)	Arrow-head	
65	25	5	5	5	5	++
	12.5	5	5	5	5	+
	6.25	5	5	5	5	+
67	25	5	5	5	5	+
	12.5	5	5	5	5	+
	6.25	5	5	5	5	+
69	25	5	5	5	5	+
	12.5	5	5	5	5	+
	6.25	5	5	5	5	-
70	25	5	5	5	5	+
	12.5	5	5	5	5	+
	6.25	5	5	5	5	+
73	25	5	5	5	5	+
	12.5	5	5	5	5	-
	6.25	4.5	5	5	5	-
74	25	5	5	5	5	+
	12.5	5	5	5	5	+
	6.25	5	5	5	5	-
75	25	5	5	5	5	++
	12.5	5	5	5	5	+
	6.25	5	5	5	5	-
77	25	5	5	5	5	-
	12.5	2	4	4	3	-
	6.25	2	4	4	2	-
81	25	5	5	5	5	++
	12.5	5	5	5	5	++
	6.25	5	5	5	5	-
82	25	5	5	5	5	++
	12.5	5	5	5	5	+
	6.25	5	5	5	5	-
83	25	5	5	5	5	+
	12.5	5	5	5	5	+
	6.25	5	5	5	5	-
Control A	25	0	0	0	0	-
	12.5	0	0	0	0	-
	6.25	0	0	0	0	-

As seen in the results of Test examples 1 and 2, the compounds of the present invention showed remarkable herbicidal effect against the principal annual and perennial weeds in paddy fields in pre- and post emergence treatment.

Furthermore, it was found that the compound of the present invention showed only little phytotoxicity in pre- and post transplantation treatment.

Then the Test examples in field are shown as follows.

Test Example 3: Pre-emergence soil surface treatment

A fixed amount of field soil was filled in a round plastic case 8 cm across and 8 cm deep, and a fixed amount of seeds of crabgrass, foxtail, pigweed, buckwheat was sown followed by covering them with soil 0.5 to 1 cm thick. Then immediately a diluted solution of the compound of the present invention was applied to treat the whole surface of soil in case at a rate of 12.5 to 25 g of the compound of the present invention per are.

After the treatment the cultivation was done in a greenhouse and the herbicidal activity was observed on the 20th day. The test was carried out on 2-replication system and each average value was sought. The judging standard of the results is the same with Test Example 1. The test results are shown in Table 5.

Table 5

0104484


Test Example 3: Pre-emergence soil surface treatment

Compound No.	Dosage g/a	Herbicidal Effect			
		Foxtail	Crabgrass	Pigweed	Buckwheat
1	25 12.5	5 5	5 5	5 5	5 5
2	25 12.5	5 4.5	5 4.5	5 5	5 5
5	25 12.5	5 5	5 5	5 5	5 5
8	25 12.5	5 5	5 5	5 5	5 5
11	25 12.5	5 5	5 5	5 5	5 5
12	25 12.5	5 5	5 5	5 5	5 5
13	25 12.5	5 5	5 5	5 5	5 5
24	25 12.5	5 5	5 5	5 5	5 5
26	25 12.5	5 5	5 5	5 5	5 5
30	25 12.5	5 5	5 4.5	5 5	5 5
34	25 12.5	5 5	5 5	5 5	5 5
42	25 12.5	4 3	4.5 3	5 5	5 5
43	25 12.5	4.5 3	5 3	5 5	5 5
44	25 12.5	4.5 3	5 4	5 5	5 5

0104484

Compound No.	Dosage g/a	Herbicidal Effect			
		Foxtail	Crabgrass	Pigweed	Buckwheat
60	25	5	5	5	5
	12.5	4	3	5	5
62	25	5	5	5	5
	12.5	5	5	5	5
63	25	4	4.5	5	5
	12.5	3	4	5	5
64	25	5	5	5	5
	12.5	5	5	5	5
68	25	5	5	5	5
	12.5	5	5	5	5
71	25	5	5	5	5
	12.5	4	5	5	5
72	25	4	5	5	5
	12.5	3	5	5	5
75	25	4	4	5	5
	12.5	3	3	5	5
81	25	4	5	5	5
	12.5	3	4.5	5	5
86	25	4	5	5	5
	12.5	3	4	5	5
Control A	25	0	0	0	0
	12.5	0	0	0	0
Control B	25	4	4.5	5	5
	12.5	2	3	5	5

Control B



Test Example 4: Post-emergence treatment

0104484

A fixed amount of field soil was filled in a round plastic case 8 cm across and 8 cm deep, and a fixed amount of seeds of foxtail (or crabgrass), pigweed was sown. When they grew up to 3 to 4-leaf stage, a wettable powder containing the compound of the present invention was sprayed on the body of plants after diluting it at a rate of 12.5, 25 or 50 g of active ingredient per are.

The test was conducted on 2-replication system. Twenty days after the treatment the test results were observed on the same judging standard and the results are shown in Table 6.

Table 6

0104484

Test Example 4: Post-emergence treatment

Compound No.	Dosage g/a	Herbicidal effect		Compound No.	Dosage g/a	Herbicidal effect	
		foxtail	pigweed			foxtail	pigweed
1	50	5	5	24	50	5	5
	25	5	5		25	5	5
	12.5	4	5		12.5	5	5
2	50	5	5	26	50	5	5
	25	5	5		25	5	5
	12.5	4.5	5		12.5	5	5
5	50	5	5	30	50	5	5
	25	5	5		25	4.5	5
	12.5	5	5		12.5	3.5	5
7	50	5	5	33	50	4	5
	25	5	5		25	2.5	4.5
	12.5	4	5		12.5	2	4
11	50	5	5	35	50	5	5
	25	5	5		25	5	5
	12.5	3	5		12.5	5	5
12	50	5	5	Control B	50	5	5
	25	5	5		25	3	5
	12.5	5	5		12.5	2	4.5
13	50	5	5	Compound No.	Dosage g/a	Herbicidal effect	
	25	5	5			crabgrass	pigweed
	12.5	5	5				
16	50	5	5	41	50	5	5
	25	4	5		25	4	5
	12.5	3.5	5		12.5	3	5
20	50	5	5	45	50	4	5
	25	4	5		25	3	5
	12.5	3	5		12.5	2	5
23	50	5	5	47	50	5	5
	25	4.5	5		25	4	5
	12.5	3.5	5		12.5	3	5

0104484

Com- pound No.	Dosage g/a	Herbicidal effect		Com- pound No.	Dosage g/a	Herbicidal effect	
		crabgrass	pigweed			crabgrass	pigweed
50	50	4	5	72	50	5	5
	25	3	5		25	4	5
	12.5	2	5		12.5	3	5
57	50	5	5	74	50	5	5
	25	3	5		25	3	5
	12.5	2.5	5		12.5	2	5
59	50	5	5	76	50	5	5
	25	3.5	5		25	3	5
	12.5	2.5	5		12.5	2	5
61	50	5	5	82	50	5	5
	25	3	5		25	4	5
	12.5	2	5		12.5	3	5
65	50	5	5	84	50	5	5
	25	3	5		25	3	5
	12.5	2	5		12.5	2	5
68	50	5	5	85	50	5	5
	25	3	5		25	3	5
	12.5	2	5		12.5	2	5
71	50	5	5	Con- trol A	50	0	0
	25	3	5		25	0	0
	12.5	2	5		12.5	0	0

Test Example 5: Phytotoxicity against crops

A fixed amount of field soil was filled in a plastic vessel sized 23 cm x 4.5 cm x 12.5 cm and a fixed amount of seeds of soybean, cotton, corn, wheat, sunflower and rice was sown followed by 3-cm thick covering with soil.

Then immediately a diluted solution of the compound of the present invention was sprayed on the soil surface with a small sprayer at the rate of 25 to 50 g of the compound of the present invention.

After the treatment the crops were grown in a greenhouse and 20 days later the degree of phytotoxicity against each crop was observed. The test was carried out on 2-replication system and each average value was sought.

The judging standard of test results is the same with Test Example 2 and the results are shown in Table 7.

Table 7

Test Example: Phytotoxicity



Compound No.	Dosage g/a	Phytotoxicity against crops					
		Soybean	Cotton	Corn	Wheat	Rice	Sunflower
1	50	-	+	-	-	-	-
	25	-	-	-	-	-	-
2	50	-	+	-	-	-	-
	25	-	-	-	-	-	-
5	50	-	-	-	-	-	-
	25	-	-	-	-	-	-
8	50	-	-	-	-	-	-
	25	-	-	-	-	-	-
11	50	-	+	-	-	-	-
	25	-	-	-	-	-	-
12	50	-	-	-	-	-	-
	25	-	-	-	-	-	-
25	50	-	+	-	-	-	-
	25	-	-	-	-	-	-
30	50	-	-	-	-	-	-
	25	-	-	-	-	-	-
Control B	50	+++	++	++	+++	+	+++
	25	+	+	+	+	-	++
43	50	-	-	-	-	-	-
	25	-	-	-	-	-	-
60	50	-	-	-	-	-	-
	25	-	-	-	-	-	-
62	50	-	-	-	-	-	-
	25	-	-	-	-	-	-
64	50	-	-	-	-	-	-
	25	-	-	-	-	-	-
68	50	-	-	-	-	-	-
	25	-	-	-	-	-	-

Compound No.	Dosage g/a	Phytotoxicity against crops					
		Soybean	Cotton	Corn	Wheat	Rice	Sunflower
71	50	-	-	-	-	-	-
	25	-	-	-	-	-	-
72	50	-	-	-	-	-	-
	25	-	-	-	-	-	-
75	50	-	-	-	-	-	-
	25	-	-	-	-	-	-
81	50	-	-	-	-	-	-
	25	-	-	-	-	-	-
Control A	50	-	-	-	-	-	-
	25	-	-	-	-	-	-

As obvious from the results of Test Examples 4 and 5, the compound of the present invention proves to show very good herbicidal activity both in pre-emergence and post-emergence treatments of main weeds in the field. On the other hand, it is clear from the results of Test Example 5 that the compound of the present invention has no phytotoxicity against crops and is a suitable herbicide for use in farmlands.

WHAT WE CLAIM IS:


1. A compound of the formula:



wherein R_4 is lower alkyl, X and Y are oxygen or sulfur and n is an integer of 3 to 6;


R_1 is hydrogen or halogen; R_2 is halogen; and R_3 is hydrogen or $C_1 - C_8$ -alkyl which may have lower alkoxy.

2. A compound according to Claim 1 wherein Z is




sulfur; R_1 is chloro or fluoro; R_2 is chloro or bromo; and R_3 is $C_2 - C_4$ -alkyl.

3. A compound according to Claim 1 wherein Z is




X is oxygen or sulfur; R_1 is chloro or fluoro; R_2 is chloro or bromo; and R_3 is $C_2 - C_4$ -alkyl.


4. The compound as claimed in Claim 2 having the formula:



5. The compound as claimed in Claim 2 having the formula:




6. A herbicidal composition comprising 0.5 to 95% by weight of a compound of the formula:



wherein R_1 , R_2 , R_3 and Z are defined as in Claim 1 and 5 to 99.5% by weight of adjuvant(s).


7. A herbicidal composition according to Claim 6 wherein Z is



wherein n is 3 or 4, X and Y are


xygen or sulfur; R_1 is chloro or fluoro; R_2 is chloro or bromo; and R_3 is $C_2 - C_4$ -alkyl.

8. A herbicidal composition according to Claim 6 wherein




$C_2 - C_4$ -alkyl and n is 3 or 4; R_1 is chloro or fluoro; R_2 is chloro or bromo; and R_3 is $C_2 - C_4$ -alkyl.

9. A method for killing weeds which comprises applying to weeds or the locus thereof a herbicidally effective amount of a compound of the formula:




wherein R_1 , R_2 , R_3 and Z are defined as in Claim 1.

10. A method according to Claim 9 wherein Z is




sulfur; R_1 is chloro or fluoro; R_2 is chloro or bromo; and R_3 is $C_2 - C_4$ -alkyl.

11. A method according to Claim 9 wherein z is **0104484**




and X is oxygen or sulfur; R_1 is chloro or fluoro; R_2 is chloro or bromo; and R_3 is $\text{C}_2 - \text{C}_4$ -alkyl.

12. A process for producing a compound of the formula:




wherein R_1 is hydrogen or halogen, R_2 is halogen, R_3 is hydrogen or $\text{C}_1 - \text{C}_8$ -alkyl which may have lower alkoxy, X and Y are oxygen or sulfur and n is an integer of 3 - 6, which comprises cyclizing a compound of the formula:




wherein R_1 , R_2 , R_3 , X, Y and n are defined as above, and R_4 is lower alkyl, by heating in an inert solvent, if necessary, in the presence of an alkali.

13. A process for producing a compound of the formula:




wherein R_1 , R_2 , R_3 and n are defined as above, which comprises reacting a compound of the formula:




wherein R_1 , R_2 , R_3 and n are defined as above, with a phosphorus polysulfide.


14. A process for producing a compound of the formula:



wherein R_1 , R_2 , R_3 , X and n are defined as above, and R_4 is lower alkyl, which comprises reacting a compound of the formula:



wherein R_4 and n are defined as above, with a compound of the formula:



wherein R_1 , R_2 , R_3 and X are defined as above.



European Patent
Office

EUROPEAN SEARCH REPORT

0104484

Application number

EP 83 10 8583

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 7)
Y	GB-A-1 492 457 (MITSUBISHI) * Whole document *	1,7	C 07 D 487/04 C 07 D 231/04 C 07 D 237/04 C 07 D 243/02 C 07 D 245/02
Y	GB-A-2 071 100 (NIPPON KAYAKU) * Whole document *	1,7	A 01 N 43/90 A 01 N 47/38 // (C 07 D 487/04
Y	GB-A-1 503 244 (MITSUBISHI) * Whole document *	1,7	C 07 D 249/00) C 07 D 237/00) (C 07 D 487/04) C 07 D 249/00) C 07 D 231/00) (C 07 D 487/04) C 07 D 249/00) C 07 D 243/00) (C 07 D 487/04) C 07 D 249/00) -/-
Y	CHEMICAL ABSTRACTS, vol. 88, no. 7, 13th February 1978, page 553, no. 50904k, Columbus, Ohio, USA & JP - A - 77 83552 (MITSUBISHI CHEMICAL INDUSTRIES CO., LTD.) 12-07-1977 * Abstract *	1,7	
P,E	EP-A-0 068 822 (ROHM & HAAS)		TECHNICAL FIELDS SEARCHED (Int. Cl. 7) C 07 D 487/00 C 07 D 231/00 C 07 D 237/00 C 07 D 245/00
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 09-12-1983	Examiner ALFARO I.
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			



European Patent
Office

EUROPEAN SEARCH REPORT

0104484

Application number

EP 83 10 8583

Page 2

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 3)
			C 07 D 245/00)
			TECHNICAL FIELDS SEARCHED (Int. Cl. 3)
The present search report has been drawn up for all claims			
Place of search		Date of completion of the search	Examiner
THE HAGUE		09-12-1983	ALEARO I.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone		T : theory or principle underlying the invention	
Y : particularly relevant if combined with another document of the same category		E : earlier patent document, but published on, or after the filing date	
A : technological background		D : document cited in the application	
O : non-written disclosure		L : document cited for other reasons	
P : intermediate document		& : member of the same patent family, corresponding document	